

The impact of age on cerebral perfusion, oxygenation and metabolism during exercise in humans

Braz, Igor D; Fisher, James P

DOI:
[10.1113/JP271081](https://doi.org/10.1113/JP271081)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):
Braz, ID & Fisher, JP 2015, 'The impact of age on cerebral perfusion, oxygenation and metabolism during exercise in humans', *The Journal of Physiology*. <https://doi.org/10.1113/JP271081>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Braz, I. D. and Fisher, J. P. (2015), The impact of age on cerebral perfusion, oxygenation and metabolism during exercise in humans. The Journal of Physiology, which has been published in final form at <http://dx.doi.org/10.1113/JP271081>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Checked March 2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The Journal of Physiology

<http://jp.msubmit.net>

JP-SR-2015-271081R1

Title: The impact of age on cerebral perfusion, oxygenation and metabolism during exercise in humans

Authors: Igor Braz
James Fisher

Author Conflict: No competing interests declared

Running Title: Age, exercise and the brain

Dual Publication: No

Funding: Coordenação de Aperfeiçoamento de Pessoal de Nível Superior
(Coordination for the Improvement of Higher Education Personnel): Igor D Braz, BEX
11588/12

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

**The impact of age on cerebral perfusion, oxygenation and metabolism during exercise
in humans**

Igor D. Braz & James P. Fisher
*School of Sport, Exercise & Rehabilitation Sciences, College of Life & Environmental
Sciences, University of Birmingham, Edgbaston, Birmingham, UK*

Running title: Age, exercise and the brain
Keywords: Exercise, cerebral circulation, ageing
TOC category: Symposium review

Please send correspondence to:
James P. Fisher, PhD
School of Sport, Exercise & Rehabilitation Sciences
College of Life & Environmental Sciences
University of Birmingham, Edgbaston,
Birmingham, B15 2TT, UK
tel: +44 (0)121 414 8011
fax: +44 (0)121 414 4121
email: j.p.fisher@bham.ac.uk

24 **Abstract**

25 Age is one of the most important risk factors for dementia and stroke. Examination of
26 the cerebral circulatory responses to acute exercise in the elderly may help to pinpoint the
27 mechanisms by which exercise training can reduce the risk of brain diseases, inform the
28 optimisation of exercise training programmes and assist with the identification of age-related
29 alterations in cerebral vascular function. During low-to-moderate intensity dynamic exercise,
30 enhanced neuronal activity is accompanied by cerebral perfusion increases of ~10-30%.
31 Beyond ~60-70% maximal oxygen uptake, cerebral metabolism remains elevated but
32 perfusion in the anterior portion of the circulation returns towards baseline, substantively
33 because of a hyperventilation-mediated reduction in the partial pressure of arterial carbon
34 dioxide (PaCO₂) and cerebral vasoconstriction. Cerebral perfusion is lower in older
35 individuals, both at rest and during incremental dynamic exercise. Nevertheless, the increase
36 in the estimated cerebral metabolic rate for oxygen and the arterial-internal jugular venous
37 differences for glucose and lactate are similar in young and older individuals exercising at the
38 same relative exercise intensities. Correction for the age-related reduction in PaCO₂ during
39 exercise by the provision of supplementary CO₂ is suggested to remove ~50% of the
40 difference in cerebral perfusion between young and older individuals. A multitude of
41 candidates could account for the remaining difference, including cerebral atrophy, enhanced
42 vasoconstrictor and blunted vasodilatory pathways. In summary, age-related reductions in
43 cerebral perfusion during exercise are partly associated with a lower PaCO₂ in exercising
44 older individuals, nevertheless the cerebral extraction of glucose, lactate and oxygen appear
45 to be preserved.

46 **Abbreviations:** CMRO₂, cerebral metabolic rate of oxygen; CO₂, carbon dioxide; CVCi,
47 cerebral vascular conductance index; MCA, middle cerebral artery; NIRS, near-infrared
48 spectroscopy; OCI, O₂-carbohydrate index; OGI, O₂-glucose index; PaCO₂, partial pressure
49 of arterial carbon dioxide; P_{ET}CO₂, partial pressure of end-tidal carbon dioxide; V_{mean}, mean
50 flow velocity; VO₂max, maximal oxygen consumption; W_{max}, maximal aerobic power.

51 **Introduction**

52 Public health interventions along with health care and education improvements over
53 several decades, have led to an increase in life expectancy and global population ageing
54 (Salomon *et al.*, 2012). Age is one of the most important risk factors for brain diseases such
55 as dementia and stroke (Sacco *et al.*, 1997; Lindsay *et al.*, 2002), but even in ‘healthy ageing’
56 brain structure and function are altered (Scahill *et al.*, 2003; Burgmans *et al.*, 2011). Grey and
57 white matter blood flow decreases by ~0.5% per year from early adulthood, and despite a
58 small increase in oxygen extraction, the cerebral metabolic rate of oxygen (CMRO₂) is also
59 decreased (Leenders *et al.*, 1990). Advancing age is associated with cerebral atrophy
60 (particularly in frontal and temporal regions), altered neural signalling, and impairments in
61 aspects of cognition (e.g., working memory and processing speed) (Martin *et al.*, 1991;
62 Jagust, 2013).

63 Regular exercise can improve cerebral perfusion (Ainslie *et al.*, 2008), memory
64 (Erickson *et al.*, 2011), mental health (Blumenthal *et al.*, 1999), and reduce age-related
65 neurodegeneration (Kramer *et al.*, 1999; Colcombe *et al.*, 2006). However, the mechanisms
66 by which exercise confers such beneficial effects remains incompletely understood.
67 Examination of the cerebral responses to acute exercise in the elderly may aid the
68 identification of age-related alterations in cerebral vascular function and inform the design of
69 exercise training programmes by helping to pinpoint the mechanisms by which exercise
70 training can improve cerebral vascular function (e.g., shear stress) and reduce the risk of
71 brain diseases (Carter *et al.*, 2014; Lucas *et al.*, 2015). A disruption in the normal cerebral
72 circulatory and metabolic responses to exercise has been implicated in fatigue (Secher *et al.*,
73 2008), therefore investigating the effects of ageing on cerebral haemodynamics may also
74 further our understanding of the mechanisms underlying the well established age-related
75 reductions in exercise tolerance (Fleg *et al.*, 1995).

In this article, we will outline the influence of age on cerebral perfusion, oxygenation and metabolism during exercise in humans, briefly review the underlying mechanisms and highlight important future directions for exploration. In most developed countries, an age of 65 years indicates the start of ‘old age’ (Roebuck, 1979). In view of the paucity of available data here we incorporate studies assessing the impact of age on the cerebral circulatory responses to exercise using an older cohort with a mean age of ≥ 57 years (Table 1).

How does cerebral blood flow respond to exercise?

The earliest direct measures of cerebral blood flow in non-anesthetized humans were made by Kety and Schmidt (1948b) using the nitrous oxide inhalation technique, and studies using this technique reported that global cerebral blood flow was unchanged during exercise (Scheinberg *et al.*, 1954; Kleinerman & Sancetta, 1955; Lambertsen *et al.*, 1959). However, a change in subject position from rest (supine) to exercise (upright), an associated change in the anatomy of cerebral drainage, confounding alterations in PaCO₂, and a reduced activation in some brain regions during exercise, may in part explain these observations (Secher *et al.*, 2008).

The administration of dissolved inert radioactive gases such as ¹³³Xenon and ⁸⁵Krypton via the common carotid artery and measurement of the emitted radiation by extracranial scintillation detectors permitted the earliest determination of regional cerebral blood flow responses to exercise (Hoedt-Rasmussen, 1965). Accordingly, Olesen (1971) observed a regional increase in perfusion of the cortical sensorimotor area corresponding to the hand during contractions with the contralateral hand. A ~10-30% increase cortical blood flow is also elicited by leg cycling, as determined with the ¹³³Xenon clearance initial slope index (Jorgensen *et al.*, 1992). This is paralleled by a comparable increase in middle cerebral artery (MCA) mean blood velocity (V_{mean}) (Jorgensen *et al.*, 1992) measured using the

transcranial Doppler technique introduced by Aaslid and colleagues (Aaslid *et al.*, 1982). In fact, leg cycling bilaterally increases V_{mean} in the MCA and anterior cerebral artery, whereas rhythmic handgrip exercise performed with the right hand principally increases the left MCA V_{mean} , and calf exercise performed with the right leg predominantly increases V_{mean} in the left anterior cerebral artery (Linkis *et al.*, 1995). Positron emission tomography and single-photon emission computed tomography have confirmed the exercise-induced regional increase in cerebral perfusion and activation in the sensorimotor and premotor regions, as well as the supplementary motor area, cerebellum and insular cortex (Williamson *et al.*, 1999; Hiura *et al.*, 2014), and highlights the coupling between regional cerebral activation and perfusion during exercise.

Duplex Doppler ultrasound can be used to quantify arterial blood flow to the brain via the extracranial vessels (internal carotid and vertebral arteries) (Schoning *et al.*, 1994), because unlike the transcranial Doppler technique, both the diameter and V_{mean} of the extracranial vessels can be determined (see *Future directions* section below). In parallel with the increase in MCA V_{mean} , blood flow in the internal carotid and vertebral arteries increases by ~17% during moderate intensity leg cycling (Hellstrom *et al.*, 1996; Sato *et al.*, 2011). However, at exercise intensities above ~60% maximal oxygen uptake (VO_2max) MCA V_{mean} and internal carotid artery flow plateau and then return toward resting levels as exercise intensity increases, whereas in contrast vertebral artery flow continues to increase up to 80% VO_2max (Sato *et al.*, 2011; Smith *et al.*, 2014).

In summary, despite early reports that global cerebral blood flow is unchanged during dynamic exercise, it is now firmly established from studies in young healthy individuals that there is an increase in cerebral perfusion at low-to-moderate exercise intensities directed mainly to the activated brain structures, whereas during exercise at higher intensities perfusion in the anterior portion of the cerebral circulation returns towards resting levels.

126

127 **How does age impact the cerebral blood flow responses to exercise?**

128 An age-related reduction in cerebral blood flow was first shown over 50 years ago
129 (Kety, 1956). However, as summarised in Table 1, limited studies have investigated the
130 influence of age on the cerebral blood flow responses to exercise in older individuals.

131 Heckmann et al., (2003) were the first to employ exercise as means of assessing the
132 influence of age on cerebral circulatory regulation in humans. A more rapid increase in MCA
133 V_{mean} was observed at the onset of leg cycling in the older group compared to their younger
134 counterparts. In contrast, pulsatility index, a surrogate marker of cerebral vascular resistance,
135 increased more rapidly at the onset of exercise in the younger group. These findings suggest
136 that cerebral circulatory regulation is delayed at exercise onset in older individuals. However,
137 the precise workloads used in this study were not reported, and since maximal aerobic fitness
138 is reduced with age (Fleg *et al.*, 1995) it is possible that the older group were exercising at a
139 higher relative workload, and this may provide an alternative explanation for the findings.

140 Figure 1 summarises the results of studies that have investigated the impact of age on
141 the cerebral perfusion responses to incremental dynamic exercise. At matched relative
142 exercise intensities of 30% and 50% heart rate reserve (equivalent to low and moderate
143 intensity exercise, respectively) a similar MCA V_{mean} has been reported in young (24 ± 3
144 years) and older middle-aged (57 ± 7 years) individuals (Fisher et al., 2008). In contrast,
145 subsequent studies examining the MCA V_{mean} during incremental leg cycling exercise to
146 exhaustion reported that MCA V_{mean} was lower in an older individuals at any of the matched
147 relative (Marsden *et al.*, 2012; Fisher *et al.*, 2013; Flück *et al.*, 2014) or absolute exercise
148 intensities (Flück *et al.*, 2014) examined. These conflicting findings may relate to the smaller
149 age difference between the groups in Fisher *et al.*, (2008) (33 years), compared to Marsden *et*
150 *al* (2012) (44 years), Fisher *et al* (2013) (44 years) and Flück *et al* (2014) (42 years). Indeed,

the normal age-related reduction in resting MCA V_{mean} was not observed in Fisher *et al.*, (2008), but was evident in the other studies (Marsden *et al.*, 2012; Fisher *et al.*, 2013; Flück *et al.*, 2014). Interestingly, no significant interaction between age and exercise intensity was observed for MCA V_{mean} by Fisher *et al.* (2013) and Flück *et al.* (2014), meaning that the magnitude of exercise-induced increase in MCA V_{mean} was not different between the young and older groups. In contrast, Marsden *et al.* (2012) observed that older adults had an attenuated increase in MCA V_{mean} at low intensity exercise (28% young vs. 15% older). The reason for these discrepancies is unclear, but may relate to the relatively small sample sizes used and methodological issues surrounding the use of using the transcranial Doppler technique as index of cerebral perfusion, as discussed in more detail below. Notably, a blunted increase in cerebral vascular conductance index (CVCi, mean arterial pressure / MCA V_{mean}) at low exercise workloads in older individuals has been noted, indicative of an attenuated cerebral vasodilatory response (Fisher *et al.*, 2013). Moreover, an enhanced cerebral vasoconstriction at moderate-to-high dynamic exercise workloads is reported in older groups (Ogoh *et al.*, 2011; Fisher *et al.*, 2013). This altered cerebral vascular response may represent a normal cerebral autoregulatory response given that blood pressure increases during exercise are typically greater in older individuals, or alternatively it may be a manifestation of a change in the balance between vasodilatory and vasoconstrictor pathways, as discussed below.

In summary, the available evidence indicates that cerebral perfusion is lower both at rest and during incremental maximal exercise with increased age, irrespective of whether absolute or relative workloads are compared. The majority of the available evidence supports the view that MCA V_{mean} increases to a similar extent in young and older individuals (~10-30%) during low-to-moderate intensity dynamic exercise.

How does age impact cerebral metabolism and oxygenation during exercise?

The mass of the brain only accounts for ~2% of body mass but remarkably CMRO₂ is ~25% (~60 mL/min) of whole-body resting oxygen consumption (Kety & Schmidt, 1946). Although there is a small contribution from anaerobic glycolysis (~10%), the oxidation of glucose is the principal mechanism by which the energy demand of the brain is met during resting wakefulness. As such, the molar ratio between the cerebral consumption of oxygen and glucose (oxygen-glucose index: OGI) is slightly lower than 6:1 (~5.7) (Siesjö, 1978). Resting CMRO₂ is reported to be reduced with ageing in some (Kety, 1956; Pantano *et al.*, 1984), but not all studies (Burns & Tyrrell, 1992). Ageing also causes a reduction in cerebral metabolic rate for glucose (Nugent *et al.*, 2014), which is estimated to decline by ~6% per decade globally with most cerebral regions affected, except for the occipital cortex and cerebellum (Petit-Taboue *et al.*, 1998).

Early reports that global cerebral blood flow was unchanged during exercise, also suggested that cerebral metabolism was unaltered (Madsen *et al.*, 1993), and in fact it was even concluded that "*during vigorous physical exercise the brain behaved as a steady-state organ with little or no change in cerebral circulation or metabolism*" (Zobl *et al.*, 1965). However, cerebral activation with tactile stimulation increases regional cerebral blood flow and CMRO₂ (Fox & Raichle, 1986), and the same appears to be true for CMRO₂ during exercise (Seifert *et al.*, 2009a; Smith *et al.*, 2014) although this has not been universally observed (Trangmar *et al.*, 2014). The cerebral metabolic rate for glucose and the OGI tend to be similar at rest and during exercise (Ide *et al.*, 2000b), although OGI can be reduced by very strenuous exercise, such as prolonged maximal exercise in the heat (Nybo *et al.*, 2003). Along with glucose, lactate also plays an important role as a substrate during exercise (Smith *et al.*, 2003), particularly when arterial lactate concentration is elevated such as during high intensity exercise. The combined uptake of glucose and lactate relative to oxygen remains

201 stable at low-to-moderate exercise intensities, but when exercise becomes more strenuous
202 glucose and lactate uptake increase in excess of oxygen, in an intensity dependent manner.
203 This means that the ‘oxygen-to-carbohydrate consumption index’ (OCI; $O_2/[glucose + \frac{1}{2}$
204 lactate]) is reduced, and during all-out rowing the OCI can decrease to <35% of the baseline
205 value (Volianitis *et al.*, 2008). As the increase in cerebral uptake of lactate does not result in
206 an accumulation of this substance in brain structures or in the cerebral spinal fluid it is
207 seemingly metabolized by the brain during exercise (Dalsgaard *et al.*, 2004).

208 To date a single study has compared the arterial–jugular venous concentration
209 differences for oxygen, glucose and lactate during exercise in young and elderly individuals
210 (Fisher *et al.*, 2013). A discontinuous incremental exercise protocol was conducted to
211 exhaustion and responses at relative workloads of 25%, 50%, 75% and 100% of maximal
212 aerobic power (W_{max}) were compared. Despite reductions in the cerebral perfusion of the
213 older group, increases estimated $CMRO_2$ and arterial–jugular venous differences for oxygen
214 (rest vs. 100% W_{max}) and lactate (rest vs. 75% and 100% W_{max}) were observed. The arterial-
215 jugular venous differences for glucose and OGI were unchanged, while the OCI was reduced
216 similarly during exercise in young and older individuals (rest vs. 75% and 100% W_{max}).
217 These findings suggest that the brain’s ability to uptake glucose and lactate is preserved with
218 healthy ageing.

219 Despite an age-related decline in maximal aerobic exercise capacity cerebral
220 oxygenation falls to a similar extent in young and older individuals during exhaustive
221 exercise, when indexed using estimates of either cerebral capillary oxygen saturation,
222 capillary oxygen tension and mitochondrial oxygen tension derived using arterial-to-jugular
223 venous differences (Fisher *et al.*, 2013), or frontal lobe oxygenation determined using near-
224 infrared spectroscopy (NIRS) (Flück *et al.*, 2014). Such observations might imply the
225 existence of a common centrally mediated element to fatigue. Rassmussen *et al.* (2010)

observed that reductions in cerebral mitochondrial oxygen tension induced by hypoxic exercise resulted in a reduction in the maximal volitional activation of the skeletal muscles (i.e., central fatigue). Intriguingly, despite maximal exercise capacity being increased by ~20% following exercise training, the reduction in cerebral mitochondrial oxygen tension at exhaustion was not different to the pre-training value (Seifert *et al.*, 2009a). Furthermore, at exhaustion the reduction in cerebral mitochondrial oxygen tension was similar before and after β -adrenergic blockade, despite blockade significantly reducing the maximal absolute workload performed (Seifert *et al.*, 2009b). However, administration of supplemental CO₂ to limit the exercise-induced fall in cerebral perfusion and oxygenation in young and older individuals fails to enhance exercise performance (Subudhi *et al.*, 2011; Flück *et al.*, 2014).

In summary, a preserved capacity for the cerebral extraction of glucose, lactate and oxygen has been observed in exercising healthy elderly individuals. Although reductions in cerebral oxygenation in young and older participants appear similar during exhaustive exercise, there is limited evidence to suggest that this limits exercise performance under normoxic conditions in healthy young and old individuals.

Age-related alterations in cerebral blood flow regulation during exercise

The regulation of cerebral blood flow in exercising humans is complex and incompletely understood, and as reviewed elsewhere, metabolic, chemical, autoregulatory, neurogenic, and systemic factors are among the likely contributors (Querido & Sheel, 2007; Secher *et al.*, 2008; Ainslie & Duffin, 2009). A brief discussion of the influence of ageing on these mechanisms follows and is summarised in Figure 2. Due to space constraints a focus will be placed on human studies where possible.

Metabolic: The molecular pathways involved in the coupling of regional cerebral activation and perfusion during exercise is incompletely understood. Among the substances

implicated in pial vessel dilatation and blood flow increases during cerebral activation are adenosine (Ko et al., 1990), the lactate/pyruvate ratio (Ido et al., 2001) and neuronal nitric oxide (Ma et al., 1996). Ageing impairs endothelial nitric oxide synthase, acetylcholine and ADP dependant cerebral vascular reactivity in aged rats (Mayhan *et al.*, 2008). In humans, alterations in several vasodilatory mechanisms (e.g., prostaglandins, ATP, nitric oxide) have been implicated in the age-related impairments in skeletal muscle vasculature regulation during exercise (Dinenno & Joyner, 2006). The use of pharmacological dissection to determine whether age-related alterations in the aforementioned molecular pathways occur within the human cerebral vasculature during exercise would greatly enhance our understanding of this topic.

Chemical: PaCO₂ is a major regulator of cerebral blood flow during exercise. PaCO₂ is generally well maintained or may increase slightly during low-to-moderate intensity exercise and may make a small contribution to the elevation of the cerebral perfusion at these workloads (Moraine *et al.*, 1993). At higher exercise intensities PaCO₂ is reduced due to hyperventilation and this restricts the exercise-induced increase in cerebral perfusion. Indeed, the provision of supplemental CO₂ during high intensity exercise in young healthy participants, in order to prevent the hyperventilation-mediated fall in PaCO₂ and maintain the partial pressure of end-tidal CO₂ (P_{ET}CO₂) at 50 mmHg, increases MCA V_{mean} by ~40% and cerebral oxygenation by ~15% (Subudhi *et al.*, 2011). Some studies have observed that elderly individuals have a reduced arterial, alveolar and P_{ET}CO₂ (Terman & Newton, 1964; Fisher *et al.*, 2013; Flück *et al.*, 2014). To assess influence of the age-related reduction in PaCO₂ on the cerebral circulatory responses to exercise, Flück *et al.* (2014) administered supplemental CO₂ to the inspired air (when PaCO₂ dropped below 40 mmHg) in order to prevent a hyperventilation-mediated reduction in PaCO₂ during incremental exhaustive leg cycling in healthy young and older individuals. Correction for the age-related difference in

276 PaCO₂ was suggested to account for ~50% of the reduction in cerebral perfusion during
277 exercise in elderly individuals. A lower cerebrovascular responsiveness to CO₂ could also
278 contribute to the lower cerebral perfusion during exercise in older individuals. However,
279 Murrell *et al.* (2013), observed that the cerebrovascular responsiveness to hypercapnia (5%
280 CO₂ added to the inspired air) was increased similarly from rest to sub-maximal exercise in
281 young and older individuals.

282 Of note, several studies mentioned above did not directly measure arterial blood
283 gases, and instead reported P_{ET}CO₂ or used P_{ET}CO₂ to calculate PaCO₂ (Fisher *et al.*, 2008;
284 Marsden *et al.*, 2012; Murrell *et al.*, 2013; Flück *et al.*, 2014). The use of P_{ET}CO₂ may lead to
285 an overestimation in changes of PaCO₂ during exercise (Robbins *et al.*, 1990; St Croix *et al.*,
286 1995), particularly when age-related changes in lung structure and function are present
287 (Miller & Tenney, 1956). However, a mathematical correction can facilitate the successful
288 estimation of PaCO₂ from P_{ET}CO₂ (Jones *et al.*, 1979) even in the elderly (St Croix *et al.*,
289 1995). A further point of note, is the observation that PaCO₂ vasodilates pial arterioles
290 principally as a consequence of local changes in extravascular pH (Kontos *et al.*, 1977a;
291 Kontos *et al.*, 1977b). Despite the observed reduction in baseline PaCO₂ with increased age,
292 arterial and arterialized pH were not altered (Fisher *et al.*, 2013; Flück *et al.*, 2014). However,
293 differences in extravascular pH, and thus a contribution to age-related alterations in cerebral
294 blood flow regulation, cannot be ruled out.

295 Along with PaCO₂, PaO₂ also modulates cerebral blood flow. The breathing of a
296 hypoxic gas evokes cerebral vasodilation *per se* (Kety & Schmidt, 1948a), however the
297 accompanying activation of the peripheral chemoreceptors causes hyperventilation, a
298 lowering of PaCO₂ and thus cerebral vasoconstriction. This phenomenon explains why during
299 acute hypoxia the exercise-induced change in MCA V_{mean} is similar to when breathing
300 normoxic air in young individuals (Ainslie *et al.*, 2007). Breathing a hyperoxic gas mixture at

rest, at least at sea level, also evokes cerebral vasoconstriction due to a chemoreflex mechanism (Floyd *et al.*, 2003). Interestingly, there is a notable regional heterogeneity in the cerebral perfusion response to hyperoxic exercise and that while changes from rest in MCA V_{mean} are unaffected a much greater response is seen in the posterior circulation (Smith *et al.*, 2012). Studies assessing the impact of hypoxia and hyperoxia on the cerebral blood flow responses to exercise in older individuals are needed to improve our understanding of the influence of arterial oxygen tension on cerebral perfusion in this population.

Blood pressure: The cerebral circulation has the intrinsic ability to maintain its flow relatively constant over a range of arterial blood pressure values (Paulson *et al.*, 1990). During dynamic exercise where pronounced intensity dependent increases in blood pressure occur, particularly in older individuals, cerebral autoregulation is likely important in order to restrict the increases in cerebral perfusion, which are modest and unlike blood pressure are greatest at low-to-moderate intensities, at least in the anterior portion of the circulation (Fisher *et al.*, 2008). Dynamic cerebral autoregulation, as determined from the linear transfer function analysis of blood pressure and MCA V_{mean} , appears to be maintained during exercise in young healthy individuals (Brys *et al.*, 2003). Similarly, the transfer function gain between mean arterial pressure and MCA V_{mean} in the very low and low frequency ranges, is not different between young and middle-aged individuals, suggesting that cerebral autoregulatory capacity is similar (Fisher *et al.*, 2008). However, the coherence between blood pressure and MCA V_{mean} noted in this study was relatively low which could have a bearing on the interpretation of these results, and carefully designed studies are required to examine whether the dynamic cerebral autoregulation of cerebral blood flow is altered in more elderly individuals during exercise.

Neurogenic: The adrenergic innervation of the cerebral vasculature has been long recognised (Lowe & Gilboe, 1971) but the nature of the sympathetic influence on the cerebral

circulation is still much debated (van Lieshout & Secher, 2008). Cerebral vasoconstriction is evoked by sympathetic nerves stimulation in animals (Auer *et al.*, 1983) and administration of α -adrenergic agonists and sympathoexcitatory manoeuvres in humans (Olesen, 1972; Micieli *et al.*, 1994). Heightened sympathetic nerve activity may serve to protect the cerebral arterioles from over-perfusion during a hypertensive insult (Bill & Linder, 1976). Therefore, a sympathetically-mediated cerebral vasoconstriction in older individuals may serve as an important mechanism to defend against the exaggerated-blood pressure response to exercise (Fisher *et al.*, 2008; Fisher *et al.*, 2013). An impairment of the normal metabolic modulation of sympathetic vasoconstrictor tone (i.e., functional sympatholysis) has been identified in the peripheral vasculature of elderly individuals (Dinenno *et al.*, 2005), although whether such a phenomenon is operative within the cerebral vasculature of elderly individuals remains to be examined. Along with the potential contribution of the sympathetic nervous system in the regulation of the cerebral blood vessels, a role for the parasympathetic system has also been suggested. Seifert *et al.* (2010) observed that the increases in MCA V_{mean} during incremental leg cycling exercise were abolished in young healthy individuals following administration of glycopyrrolate, a muscarinic cholinergic antagonist. An age-related reduction in cholinergic signalling has been identified in several tissues (e.g., heart, peripheral vasculature), and studies are needed to compare the cerebral blood flow responses to exercise in young and older participants following administration of a muscarinic cholinergic antagonist.

Central command / exercise pressor reflex: Activation of feedforward signals from brain centres, that arise in parallel with the generation of motor signals to contracting skeletal muscles (i.e., central command), evoke an accompanying increase in MCA V_{mean} (Sato *et al.*, 2009) and cerebral lactate uptake, and decrease OGI (Dalsgaard *et al.*, 2002). In addition, feedback signals from stimulation of group III and IV afferents located within the exercising skeletal muscles (i.e., exercise pressor reflex) can increase MCA V_{mean} (Braz *et al.*, 2014),

increase cerebral lactate uptake and decrease OGI (Dalsgaard *et al.*, 2003). Age-related reductions in the strength of the exercise pressor reflex have been reported (Markel *et al.*, 2003) but the implication of this for cerebral perfusion and metabolism during exercise is not known.

Systemic: Attenuating the cardiac output response to exercise in young healthy individuals with the administration of a β_1 -adrenergic receptor blocker reduces the magnitude of the normal MCA V_{mean} response (Ide *et al.*, 2000a). Furthermore, patients with atrial fibrillation in whom the cardiac output response to exercise is attenuated also demonstrate an attenuated MCA V_{mean} response (Ide *et al.*, 1999). Given the normal age-related decline in cardiac output during exercise (Hagberg *et al.*, 1985), the lower MCA V_{mean} observed during maximal exercise in elderly individuals may also relate to a lower cardiac output (increase of $\approx 200\%$ young vs. $\approx 160\%$ elderly) (Fisher *et al.*, 2013).

Aerobic fitness: Murrell *et al.* (2013) examined the MCA V_{mean} responses to incremental dynamic exercise before and after 12 weeks of aerobic exercise training which increased VO_2max from 24 ± 4 to $26 \pm 4 \text{ ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$ in older individuals (63 ± 5 years). It was observed that the MCA V_{mean} responses to exercise were unchanged following training, and that cerebrovascular responsiveness to hypercapnia was augmented at rest but not during exercise. Using a cross-sectional study design, Flück *et al.* (2014) also observed that cardiorespiratory fitness did not influence on the MCA V_{mean} responses to incremental dynamic exercise in either young (VO_2max of trained vs. untrained; 66 ± 1 vs. $50 \pm 2 \text{ ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$) or older (41 ± 3 vs. $30 \pm 1 \text{ ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$) individuals. Seifert *et al.* (2009a) showed that 3 months of aerobic exercise training attenuates the normal reduction in OCI and the increase in CMRO_2 at a matched sub-maximal exercise workload. Whether exercise training impacts the cerebral metabolic responses to exercise in elderly individuals has not been examined.

Sex: Ovarian hormones can evoke a multitude of cerebrovascular effects (Duckles & Krause, 2007). Oestrogen stimulates the production of prostacyclin and nitric oxide (Krause *et al.*, 2006), and increases MCA V_{mean} in women undergoing hormone replacement therapy (Bain *et al.*, 2004). In the three studies of the MCA V_{mean} responses to exercise that have studied both young and older, men and women no significant interactions between age and sex were noted (Table 1) (Heckmann *et al.*, 2003; Fisher *et al.*, 2008; Murrell *et al.*, 2013). However, these studies were not specifically designed to determine how sex and ovarian hormones impact the age-related changes in the cerebral circulatory responses to acute exercise in humans.

Future directions

To date, all of the studies that have examined the impact of age on the cerebral perfusion during exercise have relied upon the transcranial Doppler technique. This technique is amenable for use during exercise at a range of intensities, has high temporal and spatial resolution, and is relatively inexpensive. However, the major limitation of this technique is that the V_{mean} is only proportional to flow if the arterial diameter remains constant (Ainslie & Hoiland, 2014; Coverdale *et al.*, 2014; Verbree *et al.*, 2014). An alternative approach is the use of duplex Doppler ultrasound to assess blood flow in the extracranial arteries (Hellstrom *et al.*, 1996). This approach would also permit determination of whether ageing influences the distribution of volumetric blood flow through the internal carotid and vertebral arteries during incremental exercise, as has been performed in young individuals (Sato *et al.*, 2011). The combination of such cerebral blood flow measures with arterial-jugular venous blood sampling (Trangmar *et al.*, 2014) would also permit a more complete assessment of how age affects cerebral perfusion and metabolic nutrient delivery/removal during exercise than has been undertaken so far. In addition, studies employing a tracer dilution method are required

to determine the impact of age on cerebral lactate turnover, uptake and release, during exercise. Brain imaging techniques (e.g., functional magnetic resonance imaging) may provide an alternative approach to the assessment of cerebral perfusion, oxygenation and metabolism during exercise. Such approaches would also permit the normalisation of cerebral blood flow to cerebral mass and the consideration of regional responses with high spatial resolution. Cerebral atrophy and/or reduced neural activation might provide a potential explanation for the observed reduction in cerebral perfusion and maintained arterial-jugular venous difference for oxygen in exercising elderly individuals (Fisher *et al.*, 2013). Notably, the application of sophisticated brain imaging techniques during exercise is limited due to the incidence of movement artefacts during high intensity dynamic exercise.

At present there are no longitudinal studies assessing the impact of age on cerebral perfusion, oxygenation and metabolism during exercise in humans. Furthermore the cross-sectional studies that have been performed have utilized a single older cohort. To more completely elucidate how age impacts the cerebral response future studies should incorporate groups of individuals across a broader age range.

The vast majority of studies that have examined the impact of age on cerebral perfusion have utilized steady-state leg cycling. Although, the relative stability of the head position means that this is an attractive exercise mode to employ when assessing the cerebral vascular responses to exercise, the response profile is different to that observed during other exercise modes such as treadmill running (Lyngeraa *et al.*, 2013) and rowing (Pott *et al.*, 1997), where large fluctuations in blood pressure occur. Furthermore, the steady-state responses to exercise have predominantly been examined in young and older individuals, and carefully controlled studies investigating cerebrovascular kinetics at the onset and offset of exercise may provide important insights into the influence of age in cerebrovascular function.

The evidence base for the benefits of exercise training and physical activity for the prevention of cerebrovascular disease is substantial. However, our understanding of the precise mechanisms whereby exercise confers such beneficial effects remains incompletely understood. Among the plethora of candidate mechanisms the importance of shear stress has been highlighted (Bolduc *et al.*, 2013; Lucas *et al.*, 2015) and it is notable that the greatest exercise-induced increases in cerebral perfusion occur at low-to-moderate intensities of dynamic exercise.

Conclusions

Ageing is associated with a lower cerebral perfusion at rest and during exercise. Nevertheless, cerebral extraction of glucose, lactate and oxygen appears to be preserved in exercising older individuals. An age-related reduction in PaCO₂ has been estimated to accounts for ~50% of the lower cerebral perfusion during exercise, and other metabolic, chemical, autoregulatory, neurogenic and systemic mechanisms likely contribute. Studies are required to better understand cerebrovascular regulation during exercise in elderly individuals, to explore the utility of exercise in identifying age-related alterations in cerebral vascular function, and to optimise exercise-training regimes to promote cerebral vascular function in health and disease.

References

- Aaslid R, Markwalder TM & Nornes H. (1982). Noninvasive transcranial Doppler ultrasound recording of flow velocity in basal cerebral arteries. *Journal of neurosurgery* **57**, 769-774.
- Ainslie PN, Barach A, Murrell C, Hamlin M, Hellemans J & Ogoh S. (2007). Alterations in cerebral autoregulation and cerebral blood flow velocity during acute hypoxia: rest and exercise. *American journal of physiology Heart and circulatory physiology* **292**, H976-983.
- Ainslie PN, Cotter JD, George KP, Lucas S, Murrell C, Shave R, Thomas KN, Williams MJ & Atkinson G. (2008). Elevation in cerebral blood flow velocity with aerobic fitness throughout healthy human ageing. *The Journal of physiology* **586**, 4005-4010.
- Ainslie PN & Duffin J. (2009). Integration of cerebrovascular CO₂ reactivity and chemoreflex control of breathing: mechanisms of regulation, measurement, and interpretation. *American journal of physiology Regulatory, integrative and comparative physiology* **296**, R1473-1495.
- Ainslie PN & Hoiland RL. (2014). Transcranial Doppler ultrasound: valid, invalid, or both? *J Appl Physiol (1985)* **117**, 1081-1083.
- Auer LM, Edvinsson L & Johansson BB. (1983). Effect of sympathetic nerve stimulation and adrenoceptor blockade on pial arterial and venous calibre and on intracranial pressure in the cat. *Acta physiologica Scandinavica* **119**, 213-217.
- Bain CA, Walters MR, Lees KR & Lumsden MA. (2004). The effect of HRT on cerebral haemodynamics and cerebral vasomotor reactivity in post-menopausal women. *Hum Reprod* **19**, 2411-2414.
- Bill A & Linder J. (1976). Sympathetic control of cerebral blood flow in acute arterial hypertension. *Acta physiologica Scandinavica* **96**, 114-121.
- Blumenthal JA, Babyak MA, Moore KA, Craighead WE, Herman S, Khatri P, Waugh R, Napolitano MA, Forman LM, Appelbaum M, Doraiswamy PM & Krishnan KR. (1999). Effects of exercise training on older patients with major depression. *Archives of internal medicine* **159**, 2349-2356.
- Bolduc V, Thorin-Trescases N & Thorin E. (2013). Endothelium-dependent control of cerebrovascular functions through age: exercise for healthy cerebrovascular aging. *American journal of physiology Heart and circulatory physiology* **305**, H620-633.
- Braz ID, Scott C, Simpson LL, Springham EL, Tan BW, Balanos GM & Fisher JP. (2014). Influence of muscle metaboreceptor stimulation on middle cerebral artery blood velocity in humans. *Experimental physiology* **99**, 1478-1487.

487 Brys M, Brown CM, Marthol H, Franta R & Hilz MJ. (2003). Dynamic cerebral autoregulation remains
 488 stable during physical challenge in healthy persons. *American journal of physiology Heart*
 489 *and circulatory physiology* **285**, H1048-1054.

490

491 Burgmans S, Gronenschild EH, Fandakova Y, Shing YL, van Boxtel MP, Vuurman EF, Uylings HB, Jolles
 492 J & Raz N. (2011). Age differences in speed of processing are partially mediated by
 493 differences in axonal integrity. *NeuroImage* **55**, 1287-1297.

494

495 Burns A & Tyrrell P. (1992). Association of age with regional cerebral oxygen utilization: a positron
 496 emission tomography study. *Age and ageing* **21**, 316-320.

497

498 Carter HH, Spence AL, Pugh CJ, Ainslie P, Naylor LH & Green DJ. (2014). Cardiovascular responses to
 499 water immersion in humans: impact on cerebral perfusion. *American journal of physiology*
 500 *Regulatory, integrative and comparative physiology* **306**, R636-640.

501

502 Colcombe SJ, Erickson KI, Scalf PE, Kim JS, Prakash R, McAuley E, Elavsky S, Marquez DX, Hu L &
 503 Kramer AF. (2006). Aerobic exercise training increases brain volume in aging humans. *The*
 504 *journals of gerontology Series A, Biological sciences and medical sciences* **61**, 1166-1170.

505

506 Coverdale NS, Gati JS, Opalevych O, Perrotta A & Shoemaker JK. (2014). Cerebral blood flow velocity
 507 underestimates cerebral blood flow during modest hypercapnia and hypocapnia. *J Appl*
 508 *Physiol (1985)* **117**, 1090-1096.

509

510 Dalsgaard MK, Ide K, Cai Y, Quistorff B & Secher NH. (2002). The intent to exercise influences the
 511 cerebral O₂/carbohydrate uptake ratio in humans. *The Journal of physiology* **540**, 681-689.

512

513 Dalsgaard MK, Nybo L, Cai Y & Secher NH. (2003). Cerebral metabolism is influenced by muscle
 514 ischaemia during exercise in humans. *Experimental physiology* **88**, 297-302.

515

516 Dalsgaard MK, Quistorff B, Danielsen ER, Selmer C, Vogelsang T & Secher NH. (2004). A reduced
 517 cerebral metabolic ratio in exercise reflects metabolism and not accumulation of lactate
 518 within the human brain. *The Journal of physiology* **554**, 571-578.

519

520 Dinunno FA & Joyner MJ. (2006). Alpha-adrenergic control of skeletal muscle circulation at rest and
 521 during exercise in aging humans. *Microcirculation* **13**, 329-341.

522

523 Dinunno FA, Masuki S & Joyner MJ. (2005). Impaired modulation of sympathetic alpha-adrenergic
 524 vasoconstriction in contracting forearm muscle of ageing men. *The Journal of physiology*
 525 **567**, 311-321.

526

527 Duckles SP & Krause DN. (2007). Cerebrovascular effects of oestrogen: multiplicity of action. *Clinical*
 528 *and experimental pharmacology & physiology* **34**, 801-808.

529

530 Erickson KI, Voss MW, Prakash RS, Basak C, Szabo A, Chaddock L, Kim JS, Heo S, Alves H, White SM,
 531 Wojcicki TR, Mailey E, Vieira VJ, Martin SA, Pence BD, Woods JA, McAuley E & Kramer AF.
 532 (2011). Exercise training increases size of hippocampus and improves memory. *Proceedings*
 533 *of the National Academy of Sciences of the United States of America* **108**, 3017-3022.

534
 535 Fisher JP, Hartwich D, Seifert T, Olesen ND, McNulty CL, Nielsen HB, van Lieshout JJ & Secher NH.
 536 (2013). Cerebral perfusion, oxygenation and metabolism during exercise in young and
 537 elderly individuals. *The Journal of physiology* **591**, 1859-1870.

538
 539 Fisher JP, Ogoh S, Young CN, Raven PB & Fadel PJ. (2008). Regulation of middle cerebral artery blood
 540 velocity during dynamic exercise in humans: influence of aging. *J Appl Physiol (1985)* **105**,
 541 266-273.

542
 543 Fleg JL, O'Connor F, Gerstenblith G, Becker LC, Clulow J, Schulman SP & Lakatta EG. (1995). Impact of
 544 age on the cardiovascular response to dynamic upright exercise in healthy men and women.
 545 *J Appl Physiol (1985)* **78**, 890-900.

546
 547 Floyd TF, Clark JM, Gelfand R, Detre JA, Ratcliffe S, Guvakov D, Lambertsen CJ & Eckenhoof RG.
 548 (2003). Independent cerebral vasoconstrictive effects of hyperoxia and accompanying
 549 arterial hypocapnia at 1 ATA. *J Appl Physiol (1985)* **95**, 2453-2461.

550
 551 Flück D, Braz ID, Keiser S, Huppin F, Haider T, Hilty M, Fisher JP & Lundby C. (2014). Age, aerobic
 552 fitness and cerebral perfusion during exercise: role of carbon dioxide. *American journal of*
 553 *physiology Heart and circulatory physiology*.

554
 555 Fox PT & Raichle ME. (1986). Focal physiological uncoupling of cerebral blood flow and oxidative
 556 metabolism during somatosensory stimulation in human subjects. *Proceedings of the*
 557 *National Academy of Sciences of the United States of America* **83**, 1140-1144.

558
 559 Hagberg JM, Allen WK, Seals DR, Hurley BF, Ehsani AA & Holloszy JO. (1985). A hemodynamic
 560 comparison of young and older endurance athletes during exercise. *J Appl Physiol (1985)* **58**,
 561 2041-2046.

562
 563 Heckmann JG, Brown CM, Cheregi M, Hilz MJ & Neundorfer B. (2003). Delayed cerebrovascular
 564 autoregulatory response to ergometer exercise in normotensive elderly humans.
 565 *Cerebrovasc Dis* **16**, 423-429.

566
 567 Hellstrom G, Fischer-Colbrie W, Wahlgren NG & Jogestrand T. (1996). Carotid artery blood flow and
 568 middle cerebral artery blood flow velocity during physical exercise. *J Appl Physiol (1985)* **81**,
 569 413-418.

570
 571 Hiura M, Nariai T, Ishii K, Sakata M, Oda K, Toyohara J & Ishiwata K. (2014). Changes in cerebral
 572 blood flow during steady-state cycling exercise: a study using oxygen-15-labeled water with

573 PET. *Journal of cerebral blood flow and metabolism : official journal of the International*
574 *Society of Cerebral Blood Flow and Metabolism* **34**, 389-396.

575

576 Hoedt-Rasmussen K. (1965). Regional cerebral flow in man measured externally following intra-
577 arterial administration of 85-Kr or 133-Xe dissolved in saline. *Acta neurologica Scandinavica*
578 *Supplementum* **14**, 65-68.

579

580 Ide K, Boushel R, Sorensen HM, Fernandes A, Cai Y, Pott F & Secher NH. (2000a). Middle cerebral
581 artery blood velocity during exercise with beta-1 adrenergic and unilateral stellate ganglion
582 blockade in humans. *Acta physiologica Scandinavica* **170**, 33-38.

583

584 Ide K, Gullov AL, Pott F, Van Lieshout JJ, Koefoed BG, Petersen P & Secher NH. (1999). Middle
585 cerebral artery blood velocity during exercise in patients with atrial fibrillation. *Clin Physiol*
586 **19**, 284-289.

587

588 Ide K, Schmalbruch IK, Quistorff B, Horn A & Secher NH. (2000b). Lactate, glucose and O₂ uptake in
589 human brain during recovery from maximal exercise. *The Journal of physiology* **522 Pt 1**,
590 159-164.

591

592 Ido Y, Chang K, Woolsey TA & Williamson JR. (2001). NADH: sensor of blood flow need in brain,
593 muscle, and other tissues. *FASEB journal : official publication of the Federation of American*
594 *Societies for Experimental Biology* **15**, 1419-1421.

595

596 Jagust W. (2013). Vulnerable neural systems and the borderland of brain aging and
597 neurodegeneration. *Neuron* **77**, 219-234.

598

599 Jones NL, Robertson DG & Kane JW. (1979). Difference between end-tidal and arterial PCO₂ in
600 exercise. *Journal of applied physiology: respiratory, environmental and exercise physiology*
601 **47**, 954-960.

602

603 Jorgensen LG, Perko G & Secher NH. (1992). Regional cerebral artery mean flow velocity and blood
604 flow during dynamic exercise in humans. *J Appl Physiol (1985)* **73**, 1825-1830.

605

606 Kety SS. (1956). Human cerebral blood flow and oxygen consumption as related to aging. *Journal of*
607 *chronic diseases* **3**, 478-486.

608

609 Kety SS & Schmidt CF. (1946). The effects of active and passive hyperventilation on cerebral blood
610 flow, cerebral oxygen consumption, cardiac output, and blood pressure of normal young
611 men. *The Journal of clinical investigation* **25**, 107-119.

612

613 Kety SS & Schmidt CF. (1948a). The Effects of Altered Arterial Tensions of Carbon Dioxide and
614 Oxygen on Cerebral Blood Flow and Cerebral Oxygen Consumption of Normal Young Men.
615 *The Journal of clinical investigation* **27**, 484-492.

616
617 Kety SS & Schmidt CF. (1948b). The Nitrous Oxide Method for the Quantitative Determination of
618 Cerebral Blood Flow in Man: Theory, Procedure and Normal Values. *The Journal of clinical*
619 *investigation* **27**, 476-483.

620
621 Kleinerman J & Sancetta SM. (1955). Effect of mild steady state exercise on cerebral and general
622 haemodynamics of normal untrained subjects. *The Journal of clinical investigation* **34**, 1.

623
624 Ko KR, Ngai AC & Winn HR. (1990). Role of adenosine in regulation of regional cerebral blood flow in
625 sensory cortex. *The American journal of physiology* **259**, H1703-1708.

626
627 Kontos HA, Raper AJ & Patterson JL. (1977a). Analysis of vasoactivity of local pH, PCO₂ and
628 bicarbonate on pial vessels. *Stroke; a journal of cerebral circulation* **8**, 358-360.

629
630 Kontos HA, Wei EP, Raper AJ & Patterson JL, Jr. (1977b). Local mechanism of CO₂ action of cat pial
631 arterioles. *Stroke; a journal of cerebral circulation* **8**, 226-229.

632
633 Kramer AF, Hahn S, Cohen NJ, Banich MT, McAuley E, Harrison CR, Chason J, Vakil E, Bardell L,
634 Boileau RA & Colcombe A. (1999). Ageing, fitness and neurocognitive function. *Nature* **400**,
635 418-419.

636
637 Krause DN, Duckles SP & Pelligrino DA. (2006). Influence of sex steroid hormones on cerebrovascular
638 function. *J Appl Physiol (1985)* **101**, 1252-1261.

639
640 Lambertsen CJ, Owen SG, Wendel H, Stroud MW, Lurie AA, Lochner W & Clark GF. (1959).
641 *Respiratory and cerebral circulatory control during exercise at .21 and 2.0 atmospheres*
642 *inspired pO₂*, vol. 14.

643
644 Leenders KL, Perani D, Lammertsma AA, Heather JD, Buckingham P, Healy MJ, Gibbs JM, Wise RJ,
645 Hatazawa J, Herold S & et al. (1990). Cerebral blood flow, blood volume and oxygen
646 utilization. Normal values and effect of age. *Brain : a journal of neurology* **113 (Pt 1)**, 27-47.

647
648 Lindsay J, Laurin D, Verreault R, Hebert R, Helliwell B, Hill GB & McDowell I. (2002). Risk factors for
649 Alzheimer's disease: a prospective analysis from the Canadian Study of Health and Aging.
650 *American journal of epidemiology* **156**, 445-453.

651
652 Linkis P, Jorgensen LG, Olesen HL, Madsen PL, Lassen NA & Secher NH. (1995). Dynamic exercise
653 enhances regional cerebral artery mean flow velocity. *J Appl Physiol (1985)* **78**, 12-16.

654
655 Lowe RF & Gilboe DD. (1971). Demonstration of alpha and beta adrenergic receptors in canine
656 cerebral vasculature. *Stroke; a journal of cerebral circulation* **2**, 193-200.

657

658 Lucas SJ, Cotter JD, Brassard P & Bailey DM. (2015). High-intensity interval exercise and
659 cerebrovascular health: curiosity, cause, and consequence. *Journal of cerebral blood flow*
660 *and metabolism : official journal of the International Society of Cerebral Blood Flow and*
661 *Metabolism*.

662
663 Lyngeraa TS, Pedersen LM, Mantoni T, Belhage B, Rasmussen LS, van Lieshout JJ & Pott FC. (2013).
664 Middle cerebral artery blood velocity during running. *Scandinavian journal of medicine &*
665 *science in sports* **23**, e32-37.

666
667 Ma J, Ayata C, Huang PL, Fishman MC & Moskowitz MA. (1996). Regional cerebral blood flow
668 response to vibrissal stimulation in mice lacking type I NOS gene expression. *The American*
669 *journal of physiology* **270**, H1085-1090.

670
671 Madsen PL, Sperling BK, Warming T, Schmidt JF, Secher NH, Wildschiodtz G, Holm S & Lassen NA.
672 (1993). Middle cerebral artery blood velocity and cerebral blood flow and O₂ uptake during
673 dynamic exercise. *J Appl Physiol* (1985) **74**, 245-250.

674
675 Markel TA, Daley JC, 3rd, Hogeman CS, Herr MD, Khan MH, Gray KS, Kunselman AR & Sinoway LI.
676 (2003). Aging and the exercise pressor reflex in humans. *Circulation* **107**, 675-678.

677
678 Marsden KR, Haykowsky MJ, Smirl JD, Jones H, Nelson MD, Altamirano-Diaz LA, Gelinas JC, Tzeng YC,
679 Smith KJ, Willie CK, Bailey DM & Ainslie PN. (2012). Aging blunts hyperventilation-induced
680 hypocapnia and reduction in cerebral blood flow velocity during maximal exercise. *Age*
681 *(Dordr)* **34**, 725-735.

682
683 Martin AJ, Friston KJ, Colebatch JG & Frackowiak RS. (1991). Decreases in regional cerebral blood
684 flow with normal aging. *Journal of cerebral blood flow and metabolism : official journal of*
685 *the International Society of Cerebral Blood Flow and Metabolism* **11**, 684-689.

686
687 Mayhan WG, Arrick DM, Sharpe GM & Sun H. (2008). Age-related alterations in reactivity of cerebral
688 arterioles: role of oxidative stress. *Microcirculation* **15**, 225-236.

689
690 Micieli G, Tassorelli C, Bosone D, Cavallini A, Viotti E & Nappi G. (1994). Intracerebral vascular
691 changes induced by cold pressor test: a model of sympathetic activation. *Neurological*
692 *research* **16**, 163-167.

693
694 Miller RM & Tenney SM. (1956). Dead space ventilation in old age. *J Appl Physiol* **9**, 321-327.

695
696 Moraine JJ, Lamotte M, Berre J, Niset G, Leduc A & Naeije R. (1993). Relationship of middle cerebral
697 artery blood flow velocity to intensity during dynamic exercise in normal subjects. *European*
698 *journal of applied physiology and occupational physiology* **67**, 35-38.

699

700 Murrell CJ, Cotter JD, Thomas KN, Lucas SJ, Williams MJ & Ainslie PN. (2013). Cerebral blood flow
 701 and cerebrovascular reactivity at rest and during sub-maximal exercise: effect of age and 12-
 702 week exercise training. *Age (Dordr)* **35**, 905-920.

703

704 Nugent S, Tremblay S, Chen KW, Ayutyanont N, Roontiva A, Castellano CA, Fortier M, Roy M,
 705 Courchesne-Loyer A, Bockt C, Lepage M, Turcotte E, Fulop T, Reiman EM & Cunnane SC.
 706 (2014). Brain glucose and acetoacetate metabolism: a comparison of young and older adults.
 707 *Neurobiology of aging* **35**, 1386-1395.

708

709 Nybo L, Nielsen B, Blomstrand E, Moller K & Secher N. (2003). Neurohumoral responses during
 710 prolonged exercise in humans. *J Appl Physiol (1985)* **95**, 1125-1131.

711

712 Ogoh S, Fisher JP, Young CN & Fadel PJ. (2011). Impact of age on critical closing pressure of the
 713 cerebral circulation during dynamic exercise in humans. *Experimental physiology* **96**, 417-
 714 425.

715

716 Olesen J. (1971). Contralateral focal increase of cerebral blood flow in man during arm work. *Brain :
 717 a journal of neurology* **94**, 635-646.

718

719 Olesen J. (1972). The effect of intracarotid epinephrine, norepinephrine, and angiotensin on the
 720 regional cerebral blood flow in man. *Neurology* **22**, 978-987.

721

722 Pantano P, Baron JC, Lebrun-Grandie P, Duquesnoy N, Bousser MG & Comar D. (1984). Regional
 723 cerebral blood flow and oxygen consumption in human aging. *Stroke; a journal of cerebral
 724 circulation* **15**, 635-641.

725

726 Paulson OB, Strandgaard S & Edvinsson L. (1990). Cerebral autoregulation. *Cerebrovascular and
 727 brain metabolism reviews* **2**, 161-192.

728

729 Petit-Taboue MC, Landeau B, Desson JF, Desgranges B & Baron JC. (1998). Effects of healthy aging on
 730 the regional cerebral metabolic rate of glucose assessed with statistical parametric mapping.
 731 *NeuroImage* **7**, 176-184.

732

733 Pott F, Knudsen L, Nowak M, Nielsen HB, Hanel B & Secher NH. (1997). Middle cerebral artery blood
 734 velocity during rowing. *Acta physiologica Scandinavica* **160**, 251-255.

735

736 Querido JS & Sheel AW. (2007). Regulation of cerebral blood flow during exercise. *Sports Med* **37**,
 737 765-782.

738

739 Rasmussen P, Nielsen J, Overgaard M, Krogh-Madsen R, Gjedde A, Secher NH & Petersen NC. (2010).
 740 Reduced muscle activation during exercise related to brain oxygenation and metabolism in
 741 humans. *The Journal of physiology* **588**, 1985-1995.

742

743 Robbins PA, Conway J, Cunningham DA, Khamnei S & Paterson DJ. (1990). A comparison of indirect
 744 methods for continuous estimation of arterial PCO₂ in men. *J Appl Physiol (1985)* **68**, 1727-
 745 1731.

746

747 Roebuck J. (1979). When Does “Old Age Begin?”: The Evolution Of The English Definition. *Journal of*
 748 *Social History* **12**, 416-428.

749

750 Sacco RL, Benjamin EJ, Broderick JP, Dyken M, Easton JD, Feinberg WM, Goldstein LB, Gorelick PB,
 751 Howard G, Kittner SJ, Manolio TA, Whisnant JP & Wolf PA. (1997). American Heart
 752 Association Prevention Conference. IV. Prevention and Rehabilitation of Stroke. Risk factors.
 753 *Stroke; a journal of cerebral circulation* **28**, 1507-1517.

754

755 Salomon JA, Wang H, Freeman MK, Vos T, Flaxman AD, Lopez AD & Murray CJ. (2012). Healthy life
 756 expectancy for 187 countries, 1990-2010: a systematic analysis for the Global Burden
 757 Disease Study 2010. *Lancet* **380**, 2144-2162.

758

759 Sato K, Ogoh S, Hirasawa A, Oue A & Sadamoto T. (2011). The distribution of blood flow in the
 760 carotid and vertebral arteries during dynamic exercise in humans. *The Journal of physiology*
 761 **589**, 2847-2856.

762

763 Sato K, Sadamoto T, Ueda-Sasahara C, Shibuya K, Shimizu-Okuyama S, Osada T, Kamo M, Saito M &
 764 Kagaya A. (2009). Central command and the increase in middle cerebral artery blood flow
 765 velocity during static arm exercise in women. *Experimental physiology* **94**, 1132-1138.

766

767 Scahill RI, Frost C, Jenkins R, Whitwell JL, Rossor MN & Fox NC. (2003). A longitudinal study of brain
 768 volume changes in normal aging using serial registered magnetic resonance imaging.
 769 *Archives of neurology* **60**, 989-994.

770

771 Scheinberg P, Blackburn LI, Rich M & Saslaw M. (1954). Effects of vigorous physical exercise on
 772 cerebral circulation and metabolism. *The American journal of medicine* **16**, 549-554.

773

774 Schoning M, Walter J & Scheel P. (1994). Estimation of cerebral blood flow through color duplex
 775 sonography of the carotid and vertebral arteries in healthy adults. *Stroke; a journal of*
 776 *cerebral circulation* **25**, 17-22.

777

778 Secher NH, Seifert T & Van Lieshout JJ. (2008). Cerebral blood flow and metabolism during exercise:
 779 implications for fatigue. *J Appl Physiol (1985)* **104**, 306-314.

780

781 Seifert T, Fisher JP, Young CN, Hartwich D, Ogoh S, Raven PB, Fadel PJ & Secher NH. (2010).
 782 Glycopyrrolate abolishes the exercise-induced increase in cerebral perfusion in humans.
 783 *Experimental physiology* **95**, 1016-1025.

784

785 Seifert T, Rasmussen P, Brassard P, Homann PH, Wissenberg M, Nordby P, Stallknecht B, Secher NH
786 & Nielsen HB. (2009a). Cerebral oxygenation and metabolism during exercise following three
787 months of endurance training in healthy overweight males. *American journal of physiology*
788 *Regulatory, integrative and comparative physiology* **297**, R867-876.

789
790 Seifert T, Rasmussen P, Secher NH & Nielsen HB. (2009b). Cerebral oxygenation decreases during
791 exercise in humans with beta-adrenergic blockade. *Acta Physiol (Oxf)* **196**, 295-302.

792
793 Siesjö B. (1978). Brain energy metabolism. *Annals of Neurology* **5**, 308-308.

794
795 Smith D, Pernet A, Hallett WA, Bingham E, Marsden PK & Amiel SA. (2003). Lactate: a preferred fuel
796 for human brain metabolism in vivo. *Journal of cerebral blood flow and metabolism : official*
797 *journal of the International Society of Cerebral Blood Flow and Metabolism* **23**, 658-664.

798
799 Smith KJ, MacLeod D, Willie CK, Lewis NC, Hoiland RL, Ikeda K, Tymko MM, Donnelly J, Day TA,
800 MacLeod N, Lucas SJ & Ainslie PN. (2014). Influence of high altitude on cerebral blood flow
801 and fuel utilization during exercise and recovery. *The Journal of physiology* **592**, 5507-5527.

802
803 Smith KJ, Wong LE, Eves ND, Koelwyn GJ, Smirl JD, Willie CK & Ainslie PN. (2012). Regional cerebral
804 blood flow distribution during exercise: influence of oxygen. *Respiratory physiology &*
805 *neurobiology* **184**, 97-105.

806
807 St Croix CM, Cunningham DA, Kowalchuk JM, McConnell AK, Kirby AS, Scheuermann BW, Petrella RJ
808 & Paterson DH. (1995). Estimation of arterial PCO₂ in the elderly. *J Appl Physiol (1985)* **79**,
809 2086-2093.

810
811 Subudhi AW, Olin JT, Dimmen AC, Polaner DM, Kayser B & Roach RC. (2011). Does cerebral oxygen
812 delivery limit incremental exercise performance? *J Appl Physiol (1985)* **111**, 1727-1734.

813
814 Terman JW & Newton JL. (1964). Changes in Alveolar and Arterial Gas Tensions as Related to
815 Altitude and Age. *J Appl Physiol* **19**, 21-24.

816
817 Trangmar SJ, Chiesa ST, Stock CG, Kalsi KK, Secher NH & Gonzalez-Alonso J. (2014). Dehydration
818 affects cerebral blood flow but not its metabolic rate for oxygen during maximal exercise in
819 trained humans. *The Journal of physiology* **592**, 3143-3160.

820
821 van Lieshout JJ & Secher NH. (2008). Last Word on Point:Counterpoint: Sympathetic activity
822 does/does not influence cerebral blood flow. *J Appl Physiol (1985)* **105**, 1374.

823
824 Verbree J, Bronzwaer AS, Ghariq E, Versluis MJ, Daemen MJ, van Buchem MA, Dahan A, van Lieshout
825 JJ & van Osch MJ. (2014). Assessment of middle cerebral artery diameter during hypocapnia
826 and hypercapnia in humans using ultra-high-field MRI. *J Appl Physiol (1985)* **117**, 1084-1089.

827

828 Volianitis S, Fabricius-Bjerre A, Overgaard A, Stromstad M, Bjarrum M, Carlson C, Petersen NT,
829 Rasmussen P, Secher NH & Nielsen HB. (2008). The cerebral metabolic ratio is not affected
830 by oxygen availability during maximal exercise in humans. *The Journal of physiology* **586**,
831 107-112.

832
833 Williamson JW, McColl R, Mathews D, Ginsburg M & Mitchell JH. (1999). Activation of the insular
834 cortex is affected by the intensity of exercise. *J Appl Physiol (1985)* **87**, 1213-1219.

835
836 Zobl EG, Talmers FN, Christensen RC & Baer LJ. (1965). Effect of exercise on the cerebral circulation
837 and metabolism. *Journal of Applied Physiology* **20**, 1289-1293.

838

839

840

841

842 **COMPETING INTERESTS**

843 The authors have no potential conflicts of interest to disclose.

844

845 **FUNDING**

846 IDB is supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior

847 (CAPES, BEX 11588/12).

848

TABLES

Table 1. Summary of studies assessing the impact of ageing on cerebral perfusion during exercise

Study	Participants	Protocol	Measurements	Cerebral perfusion	Other findings
Heckmann <i>et al.</i> (2002)	18 young (10♀, 29±5 yr), 18 older (12♀, 66±5 yr)	3 min leg cycling to ↑BP >10% and ↑HR >25%	Beat-by-beat radial NIBP, ECG, transcutaneous pCO ₂ , MCA V _{mean} , PI	Rest: ↔MCA V _{mean} Exercise: ↑MCA V _{mean} faster in older at exercise onset.	↔pCO ₂ , ↑PI delayed at exercise onset in the elderly
Fisher <i>et al.</i> (2008)	9 young (3♀, 24±3 yr), 10 older (4♀, 57±7 yr)	Steady-state leg cycling at 30% (low intensity) and 50% (moderate intensity) of HRR	Beat-by-beat finger NIBP, ECG, estimated PaCO ₂ , MCA V _{mean} , CVCi	Rest: ↔MCA V _{mean} Exercise: ↑MCA V _{mean} similar in young and older.	↔PaCO ₂ , ↔MAP and MCA V _{mean} transfer function gain and phase in young and older
Marsden <i>et al.</i> (2012)	20 young ♂ (23±4 yr), 14 older ♂ (71±10 yr)	Incremental leg cycling to volitional exhaustion	Intermittent brachial NIBP, ECG, P _{ET} CO ₂ , ventilation, VO ₂ , MCA V _{mean}	Rest: ↓MCA V _{mean} Exercise: ↑MCA V _{mean} lower in older at ≤50% VO ₂ peak.	Age: ↓P _{ET} CO ₂ . At VO ₂ peak, ↓hyperventilatory response, no hypocapnia and ↓cerebral vasoconstriction in older.
Murrell <i>et al.</i> (2013)	10 young (5♀, 23±5 yr) and 10 older (5♀, 63±5 yr)	Leg cycling at 30% and 70% HRR. Hypo- and hypercapnic challenge. Before and after 12	VO ₂ max, Beat-by-beat finger NIBP, ECG, P _{ET} CO ₂ , MCA V _{mean} ,	Rest: ↓MCA V _{mean} Exercise: ↑MCA V _{mean} lower (absolute) or similar	↔P _{ET} CO ₂ , ↔CVRCO ₂ between groups at rest and exercise. Training:

	yr)	weeks of aerobic training	ventilation	(relative) in older. No change with training.	↑hypercapnic CVR _{CO₂} and work rate similarly in both groups
Fisher <i>et al.</i> (2013)	11 young ♂ (22±1 yr), 9 older ♂ (66±2 yr)	Discontinuous incremental leg cycling to volitional exhaustion	Intra-arterial BP, ECG, PaCO ₂ , MCA V _{mean} , CVCi, arterial-jugular venous differences of oxygen, glucose, lactate, OGI, OCI, CMRO ₂ .	Rest: ↓MCA V _{mean} Exercise: ↑MCA V _{mean} similar in young and older.	Age: ↓PaCO ₂ ; ↔cerebral uptake of glucose, lactate, oxygen, OGI, OCI and CMRO ₂ between groups.
Flück <i>et al.</i> (2014)	21 young ♂ (24±3 yr), 17 older ♂ (66±4 yr)	Incremental leg cycling to volitional exhaustion with and without supplemental CO ₂	Beat-by-beat finger NIBP, HR monitor, estimated PaCO ₂ , MCA V _{mean} , CVCi, ventilation, cerebral oxygenation	Rest: ↓MCA V _{mean} Exercise: Supplemental CO ₂ reduced age-related difference in MCA V _{mean} by ~50%.	Age: ↓PaCO ₂ , ↔hypercapnic CVR _{CO₂} . Improved cerebral oxygenation with added CO ₂ but ↔ performance.

CMRO₂, cerebral metabolic rate of oxygen; CO₂, carbon dioxide; CVCi, cerebral vascular conductance index; CVR_{CO₂}, cerebral vascular reactivity to CO₂; MCA, middle cerebral artery; NIBP, non-invasive blood pressure; O, older; OCI, O₂-carbohydrate index; OGI, O₂-glucose index; PaCO₂, partial pressure of arterial carbon dioxide; V_{mean}, mean flow velocity; VO₂max, maximal oxygen consumption; ♀, women; ♂, men. ↑ denotes increase, ↓ denotes decrease, and ↔ denotes no difference.

FIGURE LEGENDS

Abstract figure: Putative mechanisms explaining the impact of age on the cerebral blood flow (CBF) responses to exercise.

CO₂, carbon dioxide; CMRO₂, cerebral metabolic rate of oxygen; %VO₂max, maximal oxygen consumption; NS, nervous system; EPR, exercise pressor reflex.

Figure 1. Percentage change from rest in middle cerebral artery mean blood velocity (MCA V_{mean}), during exercise at low (<50% VO₂max or HRR [heart rate reserve]), moderate (50-75% VO₂max or HRR), high (75-90% VO₂max) and maximal (90-100% VO₂max) intensities in young (closed symbols) and older (open symbols) individuals.

Mean values are shown for the 5 studies that have examined the impact of age on cerebral perfusion responses to incremental dynamic exercise in humans. In instances where VO₂max was not determined, HRR has been used.

Figure 2. Summary of the mechanisms whereby age may impact the cerebral blood flow responses to exercise in humans.

Red arrows denote how age modifies the contribution of a specified mechanism to the cerebral blood flow to exercise (e.g., age reduces cardiac output during exercise which likely contributes to a lower cerebral perfusion). A ‘?’ indicates where the impact of age is presently unknown.





